



NASA CR-166,306

NASA CONTRACTOR REPORT 166306

NASA-CR-166306
19820013002

Genetic Engineering Possibilities
For CELSS: A Bibliography and Summary
of Techniques

Emmett J. Johnson

LIBRARY COPY

MAR 24 1982

LANGLEY RESEARCH CENTER
LIBRARY, NASA
HAMPTON, VIRGINIA

CONTRACT NASA P.O. A 73308B (JEP)
March 1982

NASA



NF02349

Genetic Engineering Possibilities
For CELSS: A Bibliography and Summary
of Techniques

Emmett J. Johnson
Tulane University Medical School
New Orleans, LA 70112

Prepared for
Ames Research Center
under Purchase Order P.O. A 73308B (JEP)



National Aeronautics and
Space Administration

Ames Research Center
Moffett Field, California 94035



N82-20876 #

Table of Contents

	<u>Pages</u>
Part I	
Bibliography of Techniques in Genetic Engineering -- Particularly with Plants	
A. General references	1-2
B. Nitrogen fixation -- Bacteria and bacteria-plant interactions	3
Addendum	4
C. Ti plasmid and crown gall tumors	5
Part II	
Bibliography of Plant Cell Culture Techniques	6-9
Part III	
Resume of the State of the Art in Plant-Bacterial Genetic Engineering and its Potential Application in the CELSS Program	10-18

Part I
Bibliography on
Genetic Engineering
Emphasis - Plants

A. General References

- Andrainov, V. M., M. B. Amerkhanova, and I. Vinetski 1979 Chloroplast DNA cloning in Escherichia coli. II. The properties of the recombinant plasmids bearing the EcoRI fragments of pea chloroplast DNA and the cloning of the DNA sequences with rRNA genes. Genetika 15, 1918-1924.
- Andersen, K., K. T. Shanmugam, S. T. Lim, L. N. Csonka, R. Tait, H. Hennecke, D. B. Scott, S. S. M. Hom, J. F. Haury, A. Valentine and R. C. Valentine 1980 Genetic engineering in agriculture with emphasis on nitrogen fixation. Trends in Biochem. Sci. 5, 35-39.
- Butenko, R. G. 1979 Cultivation of isolated protoplasts and hybridization of somatic plant cells. Int. Rev. Cytol. 59, 323-373.
- Childs, G., S. Levy and L. H. Kedes 1979 Rapid purification of biologically active individual histone messenger RNAs by hybridization to cloned DNA linked to cellulose. Biochemistry 9, 208-213.
- Cockerham, C. C. 1980 Random and fixed effects in plant genetics. Theor. Appl. Genet. 56, 119-131.
- Doy, C. H., P. M. Gresshoff, and B. G. Rolfe 1973 Time-course of phenotypic expression of Escherichia coli gene Z following transgenesis in haploid Lycopersicon esculentum cells. Nature New Biology 244, 90-91.
- Doy, C. H., P. M. Gresshoff and B. G. Rolfe 1973 Biological and molecular evidence for transgenesis of genes from bacteria to plant cells. Proc. Natl. Acad. Sci. U.S.A. 70, 723-726.
- Fernandez, S. M., P. F. Lurquin and C. I. Kado 1978 Incorporation and maintenance of recombinant-DNA plasmid vehicles pBR313 and pCR1 in plant protoplasts. FEBS Lett 87, 277-282.
- Gonzalez, C. F. and A. K. Vidaver 1979 Syringomycin production and holcus spot disease of maize: plasmid-associated properties in Pseudomonas syringae. Curr. Microbiol. 2, 75-80.
- Hughes, B. G., F. G. White and M. A. Smith 1977 Fate of bacterial plasmid DNA during uptake by barley protoplasts FEBS Letters 79, 80-84.
- Kapitsa, O. S., L. V. Zueva, I. Vinetski, V. T. Likhachev and I. G. Bukh 1979 Relationship between the nature of beta-galactosidase in a culture of tobacco cells and experiments with transgenesis of the lac⁺ trait of Escherichia coli. Dokl Akad. Nauk. USSR 245, 465-468.

- Kleinhofs, A. and R. Behki 1977 Prospects for plant genome modification by nonconventional methods. *Ann. Rev. Genetics* 11, 79-101.
- Lurquin, P. F. and C. I. Kado 1977 *Escherichia coli* plasmid pBR313 insertion into plant protoplasts and into their nuclei. *Molec. Gen. Genet.* 154, 113-121.
- Marx, J. L. 1979 Plants: Can they live in salt water and like it? *Science* 206, 1168-1169.
- Miller, J. E. 1978 A proposal for the genetic manipulation of plant metabolism. *J. Theor. Biol.* 74, 153-154.
- Shepard, J. F., D. Bidney and E. Shahin 1980 Potato protoplasts in crop improvement. *Science* 208, 17-24.
- Truelssen, E., K. Gausing, B. Jochimeen, P. Jergensen and K. A. Marcker 1979 Cloning of soybean leghemoglobin structural gene sequences synthesized in vitro. *Nucleic Acids Res.* 1979 6, 3061-3072.
- Valentine, R. C. 1978 Genetic blueprints for new plants. *The Sciences* 18, 10-13.
- Varsanyi-Brenner, A., J. F. Gusella, C. Keys, D. E. Housman, D. Sullivan, N. Brisson and D. P. S. Verma 1979 The organization of a nuclear DNA sequence from a higher plant: molecular cloning and characterization of soybean ribosomal DNA. *Gene* 7, 317-334.
- Weber, G. and K. G. Lark 1979 An efficient plating system for rapid isolation of mutants from plant cell suspensions. *Theor. Appl. Genet.* 55, 81-86.
- Weinand, U., C. Breuschke and G. Feix 1979 Cloning of double stranded DNAs derived from polysomal mRNA of maize endosperm: isolation and characterization of zein clones. *Nucleic Acids Res.* 6, 2707-2715.
- Wildman, S. G. and P. Kwanyuen 1979 Fraction I protein and other products from tobacco for food. *Basic Life Sci.* 11, 1-18.

B. Nitrogen Fixation - Bacteria and Bacteria-Plant Interactions

- Albrecht, S. L., R. J. Maier, F. J. Manus, S. A. Russell, D. W. Emerich and H. J. Evans 1979 Hydrogenase in Rhizobium japonicum increases nitrogen fixation by nodulated soybeans. Science 203, 1255-1257.
- Benson, D. R., D. J. Arp and R. H. Burris 1979 Cell-free nitrogenase and hydrogenase from Actinorhizal root nodules. Science 205, 688-689.
- Beringer, J. E. 1980 The development of Rhizobium genetics. J. Gen. Microbiol. 116, 1-7.
- Casadesus, J. and J. Olivares 1979 Rough and fine linkage mapping of the Rhizobium meliloti chromosome. Molec. Gen. Genet. 174, 203-209.
- Casse, F., C. Boucher, J. S. Julliot, M. Michel and J. Denarie 1979 Identification and characterization of large plasmids in Rhizobium meliloti using agarose gel electrophoresis. J. Gen. Microbiol. 113, 229-242.
- Dazzo, F. B. 1979 Lectins and "recognition" in Rhizobium-legume symbiosis. ASM News 45, 238-240.
- Gross, D. C., A. K. Vidaver and R. V. Klucas 1979 Plasmids, biological properties and efficacy of nitrogen fixation in Rhizobium japonicum stains indigenous to alkaline soils. J. Gen. Microbiol. 114, 257-266.
- Kondorosi, A., E. Vincze, A. W. B. Johnston and J. Beringer 1980 A comparison of three Rhizobium linkage maps. Molec. Gen. Genet. 178, 403-408.
- Liu, S. T. and C. I. Kado 1979 Indoleacetic acid production: A plasmid function of Agrobacterium tumefaciens C58. Biochem. Biophys. Res. Comm. 90, 171-178.
- Lorkiewicz, Z., W. Zurkowski, E. Kowalczyk and A. Gorska-Melke 1971 Mutagenesis and conjugation in Rhizobium trifolii. Acta Microbiol. Polonica 3, 101-107.
- Nuti, M. P., A. A. Lepidi, R. K. Prakash, R. A. Schilperoost and F. C. Cannon 1979 Evidence for nitrogen fixation (nif) genes on indigenous Rhizobium plasmids. Nature 282, 533-535.
- Page, W. J. and M. von Tigerstrom 1979 Optimal conditions for transformation of Azotobacter vinelandii. J. Bacteriol. 139, 1058-1061.
- Puhler, A., H. J. Burkhardt and W. Klipp 1979 Cloning of the entire region for nitrogen fixation from Klebsiella pneumoniae on a multicopy plasmid vehicle in Escherichia coli. Molec. Gen. Genet. 176, 17-24.
- Urban, J. E. 1979 Non-dividing, bacteroid-like Rhizobium trifolii: In vitro induction via nutrient enrichment. Appl. and Environ. Microbiol. 38, 1173-1178.

Part I, B -- Addendum

- Brill, W. J. 1980 Biochemical genetics of nitrogen fixation
Microbiol. Rev. 44, 449-467.
- van Berkum, P. and B. B. Bohlool 1980 Evaluation of nitrogen fixation by bacteria in association with roots of tropical grasses. Microbiol. Rev. 44, 491-517.

C. Ti plasmid and Crown Gall Tumors.

- Dhaese, P., H. De Greve, H. Decraemer, J. Schell and M. Van Montagu 1979 Rapid mapping of transposon insertion and deletion mutations in the large Ti-plasmids of Agrobacterium tumerfaciens. Nucl. Acids. Res. 7, 1837-1849.
- Ellis, J. G. and A. Kerr 1979 Arginine catabolism: A new function of both octopine and nopaline Ti-plasmids of Agrobacterium. Molec. Gen. Genet. 173, 263-269.
- Gurley, W. B., J. D. Kemp, M. J. Albert, D. W. Sutton and J. Callis 1979 Transcription of Ti plasmid-derived sequences in three octopine-type crown gall tumor lines. Proc. Natl. Acad. Sci. USA. 76, 2828-2832.
- Koekman, B. P., G. Ooms, P. M. Klapwijk and R. A. Schilperoost 1979 Genetic map of an octopine Ti-plasmid. Plasmid 2, 347-357.
- Merlo, D. J., R. C. Nutter, A. L. Montoya, D. J. Garfinkel, M. H. Drummond, M. D. Chilton, M. P. Gordon and E. W. Nester 1980 The boundaries and copy numbers of Ti plasmid T-DNA vary in crown gall tumors. Molec. Gen. Genet. 177, 637-643.
- Nester, E. W. and A. Montoya 1979 Crown Gall: A natural case of genetic engineering. ASM News 45, 283-287.
- Rapp, B. J., J. D. Kemp and F. White 1979 Isolation of a non-tumor-inducing mutant of the Ti plasmid of Agrobacterium tumefaciens strain B6. Can. J. Microbiol. 25, 291-297.
- Schell, J., M. Van Montagu, M. De Beuckeleer, M. De Block, A. Depicker, M. De Wilde, G. Engler, C. Genetello, J. P. Hernalsteens, M. Holsters, J. Seurinck, B. Silva, F. Van Vliet and R. Villarroel 1979 Interactions and DNA transfer between Agrobacterium tumefaciens, the Ti-plasmid and the plant host. Proc. R. Soc. Lond. (Biol) 204, 251-266.
- Scott, I. M., J. L. Firmin, D. N. Butcher, L. M. Searle, A. K. Sogeki, J. Eagles, J. F. March, R. Self and G. R. Fenwick 1979 Analysis of a range of crown gall and normal plant tissues for Ti plasmid-determined compounds. Molec. Gen. Genet. 176, 57-65.
- Sobota, A. E. 1979 Effect of sublethal heat injury on tumour induction and RNA synthesis in Agrobacterium tumefaciens. Microbios. 23, 115-126.
- Thomashow, M. F., R. Nutter, A. L. Montoya, M. P. Gordon and E. W. Nester 1980 Integration and organization of Ti plasmid sequences in crown gall tumors. Cell 19, 729-739.
- Yang, F., A. L. Montoya, D. J. Merlo, M. H. Drummond, M-D Chilton, E. W. Nester and M. P. Gordon 1980 Foreign DNA sequences in crown gall teratomas and their fate during the loss of the tumorous traits. Molec. Gen. Genet. 177, 707-714.
- Zambryski, P., M. Holsters, K. Kruger, A. Depicker, J. Schell, M. Van Montagu and H. M. Goodman 1980 Tumor DNA structure in plant cell transformed by A. tumefaciens. Science 209, 1385-1391.

Part II
Bibliography of Plant Cell Culture Techniques

- Aviv, D. and E. Galun 1978 Biotransformation of monoterpenes by mentha cell lines: conversion of pulegone to isomenthone. *Planta. Med.* 33, 70-77.
- Bazzaz, M. B. and C. A. Rebeiz 1978 Chloroplast culture: the chlorophyll repair potential of mature chloroplasts incubated in a simple medium. *Biochim. Biophys. Acta.* 504, 310-323.
- Benigni, R., M. Bignami, I. Camoni, A. Carere, G. Conti, R. Iachetta, G. Morpurgo and V. A. Ortali 1979 A new in vitro method for testing plant metabolism in mutagenicity studies. *J. Toxicol. Environ. Health* 5, 809-819.
- Berlin, J. and U. Mutert 1978 Evidence for distinct amino acid transport systems in cultured tobacco cells. *Z. Naturforsch* 33, 641-645.
- Berlin, J. and B. Vollmer 1979 Effects of alpha-aminooxy-beta-phenylpropionic acid on phenylalanine metabolism in p-fluorophenylalanine sensitive and resistant tobacco cells. *Z. Naturforsch* 80, 770-775.
- Berlin, J. and J. M. Widholm 1978 Amino acid uptake by amino acid analog resistant tobacco cell lines. *Z. Naturforsch* 33, 634-640.
- Berlyn, M. B. 1978 A mutational approach to the study of photorespiration. *Basic Life Sci.* 11, 153-164.
- Betz, B. and K. Hahlbrock 1979 Identity of differently-induced phenylalanine ammonia-lyases from cell suspension cultures of Petroselinum hortense. *FEBS Letters* 107, 233-236.
- Betz, B., E. Shafer and K. Hahlbrock 1978 Light-induced phenylalanine ammonia-lyase in cell-suspension cultures of Petroselinum hortense. Quantitative comparison of rates of synthesis and degradation. *Arch. Biochem. Biophys* 190, 126-135.
- Chen, C. M. and D. K. Melitz 1979 Cytokinin biosynthesis in a cell-free system from cytokinin-autotrophic tobacco tissue cultures. *FEBS Letters* 107, 15-20.
- Christianson, M. L. and M. O. Chiscon 1978 Use of haploid plants as bioassay for mutagens. *Environ. Health Perspect.* 27, 77-83.
- Dixon, R. A. and C. J. Lamb 1979 Stimulation of de novo synthesis of L-phenylalanine ammonia-lyase in relation to phytoalexin accumulation in Colletotrichum lindemuthianum elicitor-treated cell suspension cultures of french bean (Phaseolus vulgaris). *Biochim. Biophys. Acta.* 586, 453-463.
- Dougall, D. K. 1979 Production of biologicals by plant cell cultures. *Adv. Exp. Med. Biol.* 118, 135-152.

- Fosket, D. E. and D. A. Tapfer 1978 Hormonal regulation of growth in cultured plant cells. *In Vitro* 14, 63-75.
- Haimer, Y. M., Y. Mizrahi and U. Bachrach 1979 Ornithine decarboxylase activity in rapidly proliferating plant cells. *FEBS Letters* 104, 146-148.
- Heins, M., J. Wahl, H. Lerch, F. Kaiser and E. Reinhard 1978 Preparation of beta-methyldigoxin by hydroxylation of beta-methyldigitoxin in fermenter cultures of *Digitalis ianata*. *Planta. Med.* 33, 57-62.
- Kemp, J. D., D. W. Sutton and E. Hack 1979 Purification and characterization of the crown gall specific enzyme nopaline synthase. *Biochemistry* 18, 3755-3760.
- Kerbaux, G. B. and K. G. Hell 1979 Effect of gamma radiation on the in vitro growth of excised pith cells of *Nicotiana tabacum* L. *Int. J. Radiat. Biol.* 35, 273-276.
- Kuboi T. and Y. Yamada 1978 Regulation of the enzyme activities related to lignin synthesis in cell aggregates of tobacco cell culture. *Biochim. Biophys. Acta* 542, 181-190.
- Kurkdjian, A., J. J. Legauy and J. Guern 1978 Measurement of intracellular pH and aspects of its control in higher plant cells cultivated in liquid medium. *Respir. Physiol.* 33, 75-89.
- Link, G., G. H. Kidd, G. Richter and L. Bogorad 1978 Structural relationships among the multiple forms of DNA-dependent RNA polymerase II from cultured parsley cells. *Eur. J. Biochem.* 91, 363-368.
- Lorz, H., W. Wernicke and I. Potrykus 1979 Culture and plant regeneration of *Hyoscyamus* protoplasts. *Planta. Med.* 36, 21-29.
- Miassod, R. and J. P. Cecchini 1979 Partial base-methylation and other structural differences in the 17 S ribosomal RNA of sycamore cells during growth in cell culture. *Biochim. Biophys. Acta.* 562, 292-301.
- Nagrutiu, I., A. Cattoir-Reynaerts and M. Jacobs 1978 Selection and characterization of cell lines of *Arabidopsis thaliana* resistant to amino acid analogs (proceedings) *Arch. Int. Physiol. Biochim* 86, 442-443.
- Olsnes, S., K. Sandvig, K. Eiklid and A. Pihl 1978 Properties and action mechanism of the toxic lectin modeccin: interaction with cell lines resistant to modeccin, abrin, and ricin. *J. Supramol. Struct.* 9, 15-25.
- Polacco, J. C. and E. A. Havir 1979 Comparisons of soybean urease isolated from seed and tissue culture. *J. Biol. Chem.* 254, 1707-1715.
- Reynolds, J. F. and T. Murashige 1979 Plant cell lines. *Methods Enzymol.* 58, 478-486.
- Ruesink, A. W. 1979 Fusion of higher plant protoplasts. *Methods Enzymol.* 58, 359-367.
- Sawai, Y., N. Sugano and K. Tsukada 1978 Ribonuclease H activity in cultured plant cells. *Biochim. Biophys. Acta.* 518, 181-185.

- Sawai, Y., S. Uchida, J. Saito, N. Sugano and K. Tsukada 1979 Two ribonucleases H from cultured plant cells. *J. Biochem. Tokyo* 85, 1301-1308.
- Schallenberg, J. and J. Berlin 1979 5-Methyltryptophan resistant cells of Catharanthus roseus. *Z. Naturforsch* 34, 541-545.
- Schrodar J., F. Kreuzaler, E. Schafer and K. Hahlbrock 1979 Concomitant induction of phenylalanine ammonia-lyase and flavanone synthase mRNAs in irradiated plant cells. *J. Biol. Chem.* 254, 57-65.
- Sharma, S. K. and S. A. Brown 1979 Affinity chromatography of Ruta graveolens L. O-methyltransferases. Studies demonstrating the potential of the technique in the mechanistic investigation of O-methyltransferases. *Can. J. Biochem.* 57, 986-994.
- Shininger, T. L. 1978 Hormone regulation of development of plant cells. *In vitro* 14, 63-75.
- Silflow, C. D. and J. L. Key 1979 Stability of polysome-associated polyadenylated RNA from soybean suspension culture cells. *Biochemistry* 18, 1013-1018.
- Silflow, C. D., J. R. Hammett and J. L. Key 1979 Sequence complexity of polyadenylated ribonucleic acid from soybean suspension culture cells. *Biochemistry* 18, 2725-2731.
- Sparkuhl, J. and G. Setterfield 1978 Vectorial transport of proteins by membrane-bound ribosomes of nongrowing and growing Jerusalem artichoke tuber cells. *Can. J. Biochem.* 56, 167-173.
- Stanley, P. and J. P. Carver 1978 Binding of (125I) wheat germ agglutinin to Chinese hamster ovary cells under conditions which affect the mobility of membrane components. *J. Cell Biol.* 79, 617-622.
- Tsang, Y. F. and R. K. Ibrahim 1979 Evidence for the existence of meta and para directing O-methyltransferases in tobacco cell cultures. *Z. Naturforsch* 34, 46-50.
- Turner, T. D. and M. H. Williams 1978 Techniques in plant cell and dispersion culture (proceeding) *J. Pharm. Pharmacol.* 30, suppl: 94 p.
- Walter, T. J., J. A. Connelly, B. G. Gengenbach and F. Wold 1979 Isolation and characterization of two homoserine dehydrogenases from maize suspension cultures. *J. Biol. Chem.* 254, 1349-1355.
- Warry, P. A. and K. M. Stoffelsen 1979 The effect of ionizing radiation on the survival of free plant cells cultivated in suspension cultures. *Int. J. Radiat. Biol.* 35, 293-298.
- Wider de Xifra, E. A., A. M. Del C. Batlle 1978 Porphyrin biosynthesis: immobilized enzymes and ligands. VI. Studies on succinyl CoA synthetase from cultured soybean cells. *Biochim. Biophys. Acta.* 523, 245-249.

- Wink, M. and T. Hartmann 1979 Cadaverine--pyruvate transamination: The principal step of enzymatic quinolizidine alkaloid biosynthesis in Lupinus polyphyllus cell suspension cultures. FEBS Letters 101, 343-346.
- Withers, L. A. 1978 A freeze-preservation of synchronously dividing cultured cells of Acer pseudonlatanus L. Cryobiology 15, 87-92.
- Zryd, J. P. 1979 Colchicine-induced resistance to antibiotic and amino-acid analogue in plant cell cultures. Experientia 35, 1168-1169.

Note: An important major source reference is the following

Perspectives in plant cell and tissue cultures. Int. Rev. Cytol. Suppl. 11A Ed. I. K. Vasil 1980 250 pps.

Part III
'Resume' of the State of the Art in Plant-Bacterial Genetic
Engineering and its Potential Application in the CELSS Program

STATEMENT OF WORK

It is a goal of the CELSS (Controlled Ecological Life Support System) Program to develop the capability to recycle materials within a closed system containing people. To accomplish this, it will be necessary to grow food on nutrients generated within the system. Such nutrients would come from wastes generated by humans and by other biological components in the system. The nutrients would be used to grow some primary producer, that is, one or a group of photosynthetic organisms.

One group of photosynthetic organisms that are of primary interest are the higher plants. To a significant extent, the characteristics of the plants within the system will determine the general size of the system, and consequently the weight and volume of such a system. Because it is anticipated that such a closed system will serve to support people in space, it is desirable to decrease the ratio of the plants - to - man volume and weight. Two ways in which this could be accomplished would be to increase the productivity of the plants, and to simplify their nutrient requirements.

For the purposes of CELSS, it is of interest to investigate the potentialities of genetic engineering of plants. The goals of such engineering might be directed toward increased productivity, or toward the development of mechanisms for nitrogen fixation within the organisms. It is also conceivable that genetic manipulation might alter the ratio or the quality of plant products.

The contractor is requested to provide the background information necessary to allow decisions on whether extensive investigation of genetic

engineering should be considered in the context of CELSS.

Given the stated goals of the CELSS program as expressed above, the following condensed review summarizes succinctly the state of the art with respect to genetic engineering possibilities particularly in regard to plant and bacteria-plant interactions.

It would seem that the following considerations are generally regarded as high priority, and any genetic engineering approaches to plants as food, whatever the system, including closed biological systems, must include them.

I. Plants as food sources - some of the essentials.

a) N₂ fixation

Nitrogen is of course a major requirement of all plants. The usual crop plants can be divided into two major classes. The cereal crops such as rice, wheat, corn, etc. require fixed forms of nitrogen such as NH_4^+ , or NO_3^- and cannot utilize atmospheric nitrogen. The other class, legumes, such as soybeans, pea and clover, to name only a few, are capable of utilizing atmospheric nitrogen (N_2) by virtue of a symbiotic relationship with certain bacteria which form root nodules on these plants. The genes responsible for this are the nif genes (for nitrogen fixing genes), a cluster known to contain at least 14 genes. The cluster can be transferred to a plasmid, and therefore is mobilizable and transferable.

There are considerations beyond N_2 fixation however, which deal with maximum efficiency of the process, and must be considered in the overall genetic engineering decisions.

Some of these considerations are as follows.

1) Energy demand of the process of N₂-fixation:

An important consideration here is the hut genes, involved in enhancing the energy efficiency of N₂-fixation. The mobilization on plasmids for transferability has been achieved.

b) CO₂-fixation

This is of course an essential event in the metabolism of all plants and photosynthetic bacteria. The responsible genes are the cfx genes. There is no reason to believe they could not be mobilized although, as yet, they have not been engineered in that way.

c) Water-splitting enzymes for generation of reductant

These are the lit genes and are critical to production of reducing power which together with ATP fulfill the energy requirements of photosynthesis.

Of lesser importance, but significant, are such things as the osm genes for osmoregulation, the cut genes for cellulose utilization, and the den genes for denitrification. All of the aforementioned genes are discussed in detail by Andersen et. al (1980 see bibliography Part I).

There is no reason to believe the mobilization and transfer of these sets of genes by plasmid vectors is not realizable. The general technology is well worked out. The specifics in each case must be separately analyzed, and successful realization of the suggested potential may require a substantial commitment in brainpower, manpower and financial support.

2) Additional considerations in reference to plants as food source in closed biological systems

a) ratio of edible to non-edible components

To my knowledge, there is no directly applicable available information with regard to genetic engineering of mobilizable and transferable genetic units for plant to plant or bacteria to plant exchanges.

C. Rate of growth

Although there doesn't appear to be any specific information on mobilization of genes involved in the rate of growth of the major crop plants, there is some information on 2 major influences in this regard. Plants seem to fall into two groups with respect to the first product of photosynthesis, the C_3 group in which 3-phospho-D-glycerate is the first product of photosynthesis and the C_4 group in which malate and aspartate are the first products of photosynthesis. More importantly, however, leaves of plants of the C_4 group have high rates of net CO_2 assimilation (42 to 85 mg CO_2 per decimeter² per hour) and low rates of photorespiration at higher levels of irradiation in normal air at 25 to 35°C. (I. Zelitch Photosynthesis and plant productivity. C & E News, Feb. 5, 1979 p. 28-48). On the other hand, with few exceptions, leaves of C_3 crop species assimilate CO_2 at rates about half or less those of C_4 plants and have rapid rates of photorespiration. The exceptions to this rule, such as sunflower and cattail, are C_3 species with rapid rates of net photosynthesis. An understanding of the way in which these unusual species overcome the handicap of rapid photorespiration exhibited by all members of the C_3 group and the mobilization and transfer of these traits would be an invaluable advantage in improving photosynthetic efficiency.

The C_4 crop species, such as sugar cane, corn, sorghum, millet and certain weeds, have average yields and average seasonal crop growth rates

(as measured by the dry weight produced per square meter of land area per week) that are two or three times greater than do C_3 species, such as wheat, soybean, spinach, tobacco and hay.

d) Altering constituents

There is to my knowledge no specific information on such things as genetic engineering to increase the protein content of potatoes or the starch content of soybeans. There is however information on recent attempts to improve potatoes by genetic engineering of potato protoplasts (Shepard, J. F., D. Bidney, and E. Shahin 1980 Potato protoplasts in crop improvement. *Science* 208, 17-24). The emphasis in this work was on resistance to disease, but other information which derived from this research is important to any considerations of genetic engineering including of course the possibility of increasing the protein content of potatoes.

In fact, by their single-cell nature and the relative ease with which they may be cultured, plant mesophyll cell protoplasts have simplistically been likened to microbes as objects for genetic manipulation. In theory, it is necessary only to induce a desired mutation within a protoplast population, allow ample time for expression, and then apply appropriate selection pressure to recover first a cell and, ultimately, a plant having a specified modification. Such a course of events would be possible, however, only if the cell is a true haploid, which is rare in plants, and the trait is simply inherited.

An additional finding with potatoes was that apart from the practical considerations the mere appearance of high frequency phenotypic variation for the several traits which were examined in "Russet Burbank" protoclonal raised the interesting genetic question of the source of such variation particularly in the absence of intentional mutagenesis. At present, the matter is purely speculative, but it is pertinent that many vegetatively propagated plants express spontaneous mutations directly and often. Among potato plants in the field, the frequency has been placed at between 1.5×10^{-3} and 1.2×10^{-5} , depending on the character and the variety (Keiken, A. Spontaneous and x-ray induced somatic aberrations in Solanum tuberosum L. Almquist and Wikell, Stockholm, 1960). Sweet potatoes (Ipomoea batatas) are capable of even higher frequencies. In the 'Centennial' cultivar, for example, two plants per 100 differed in some morphological feature. (Hammett, H. L. 1979 Hort Science 14, 123). If genetic variation is so rampant, it would seem that with the right selection pressure the desired type should be relatively easy to obtain.

e) Other considerations

1. Plant Growth Regulators

The enormous potential of plant growth regulators to growth behavior manipulation cannot be ignored. For practical purposes plant growth regulators can be defined as either natural or synthetic compounds that are applied directly to a plant to alter its life processes or structure in some

beneficial way so as to enhance yields, improve quality or facilitate harvesting. Included in this group are plant hormones (phytohormones) and herbicides when applied to induce a specific beneficial change. The four classes of hormones are auxins, giberellins, cytokinins, and inhibitors. It has been stated that giberellins probably cause changes at the gene level that result in the synthesis of new enzymes and that cytokinins also probably act at the gene level, their exact mechanism of action, however, being unknown (Nickell, L. G. 1978 Plant growth regulators - controlling biological behavior with chemicals. Chem. Eng. News, Oct. 9, 18-34). The following list of uses of plant growth regulants emphasizes their potential in the field of plant genetic and growth manipulation.

Promote rooting and propagation of plants

Initiate or terminate the dormancy of seeds, buds
and tubers

Promote or delay flowering

Induce or prevent leaf and/or fruit drop (abscission)

Control fruit set and further fruit development

Control plant or organ size

Prune the plant chemically

Increase plant resistance to pests

Enhance plant resistance to such environmental

factors as temperature, water, and air pollution

Prevent postharvest spoilage

Regulate the chemical composition of plants and
the color of fruit

Influence mineral uptake from the soil

Change the timing of crop development

2) Broad spectrum nitrogen fixing root nodule formation

One last aspect which perhaps bears separate attention is the phenomenon referred to as "faithful" and "promiscuous" nitrogen fixing root nodule formation. It is recognized that nodulation genes are a composite of contributions from the symbiotic nitrogen-fixing bacteria and the host plant. Additionally it is known that some strains of Rhizobium that form nodules on leguminous plants are very selective in their choice of host, forming nodules only on a very narrow range of plants and are therefore called "faithful". In contrast, Rhizobium such as the "cowpea" of the tropics, are said to be "promiscuous", because they form nodules on many different plants including cowpea, soybean, peanut and mung bean (see Valentine, R. C. Part I). More importantly, it has now been shown that Rhizobium can form nodules on non-leguminous plants (ibid), and that there are nitrogen fixing trees, the bacterial symbiont of which is an unusual organism called Frankia which resembles soil actinomycetes, but has not yet been cultured in the laboratory. Frankia forms nodules on a variety of woody species and thus is regarded as having evolved a relatively promiscuous set of host selection genes. Frankia nodules provide fixed nitrogen to large portions of the dry

eastern slope of the Sierra Nevada containing a mixed stand of Purshia (bitterbrush). Bitterbrush provides a high protein browse for deer and range animals. It is not known whether Frankia represents a diverse or closely related group of bacteria. Certainly, these bacteria possess a unique set of host-selectivity genes which might be harnessed for nodule formation in a wide variety of plant systems (Ibid).

The blue-green algae represent the most promiscuous symbionts found in nature since they work effectively with a variety of plants forming leaf nodules, root nodules, and trunk cavities on different plants, respectively. Any success in the genetic engineering of these versatile microorganisms would be invaluable to the kinds of efforts being discussed in this report.

Although the last word is far from being heard in genetic engineering of plant and bacteria-plant systems particularly with reference to "Controlled Ecological Life Support Systems", the collective body of information assembled in this report should allow rationale decisions about feasibility, and directional priorities.

1. Report No. NASA CR-166306		2. Government Accession No.		3. Recipient's Catalog No.	
4. Title and Subtitle Genetic Engineering Possibilities for CELSS: A Bibliography and Summary of Techniques				5. Report Date March 1982	
				6. Performing Organization Code	
7. Author(s) Emmett J. Johnson				8. Performing Organization Report No.	
9. Performing Organization Name and Address Tulane University Medical School New Orleans, LA 70112				10. Work Unit No.	
				11. Contract or Grant No. P.O. A 73308B (JEP)	
12. Sponsoring Agency Name and Address National Aeronautics and Space Administration Washington, DC 24056				13. Type of Report and Period Covered Final Report	
				14. Sponsoring Agency Code	
15. Supplementary Notes Robert D. MacElroy, Contract Monitor, Mail Stop 239-10, NASA Ames Research Center, Moffett Field, CA 94035 (415) 965-5573 FTS 448-5573 . The first in a series of CELSS reports					
16. Abstract This report provides a bibliography of the most useful techniques employed in genetic engineering of higher plants, bacteria associated with plants, and plant cell cultures. A resume of state-of-the-art genetic engineering of plants and bacteria is presented. The potential application of plant-bacterial genetic engineering to CELSS (Controlled Ecological Life Support System) program and future research needs are discussed.					
17. Key Words (Suggested by Author(s)) CELSS Genetic Engineering Life Support System Nitrogen Fixation				18. Distribution Statement Unclassified - Unlimited STAR Category - 54	
19. Security Classif. (of this report) Unclassified		20. Security Classif. (of this page) Unclassified		21. No. of Pages 21	
22. Price*					

End of Document